

Eucalyptus globulus subsp. *pseudoglobulus*

Victorian eurabbie

Classification

Eucalyptus | Symphyomyrtus | Maidenaria | Euryotae | Globulares | Euglobulares

Nomenclature

Eucalyptus globulus subsp. **pseudoglobulus** (Naudin) J.B.Kirkp., *J. Linn. Soc., Bot.* 69: 101 (1975).

Eucalyptus globulus var. *pseudoglobulus* Naudin, *Descr. Emploi Eucalyptus introd. Europe 2nd Memoire*: 34 (1891); *E. pseudoglobulus* (Naudin) Maiden, *Crit. Revis. Eucalyptus* 8: 28 (1929). T: France. Villa Thuret, *Naudin s.n.*, undated; Lectotype: P; *fide* Bean, A.R., *Telopea* 12(4): 516 (2010).

Eucalyptus globulus var. *stjohnii* R.T. Baker, *Victorian Naturalist* 30: 127 (1913); *E. stjohnii* (R.T.Baker) R.T.Baker, *Hardwoods Australia* 218 (1919). T: Lerderderg R., Vic., 5 Nov. 1903, *P.R.H.St John s.n.*; holotype: MEL [MEL1057322A].

Description

Tree to 45 m tall. Forming a lignotuber.

Bark smooth apart from base which has persistent slabs, shedding in large strips and slabs; smooth bark white, cream, grey, pink, yellowish or pale creamy orange, often with ribbons of decorticated bark in the upper branches; branchlets occasionally glaucous.

Juvenile growth (coppice or field seedlings to 50 cm): stem square in cross-section and prominently winged, glaucous; juvenile leaves opposite and sessile for many pairs, elliptical to oblong or ovate, 3.3–10 cm long, 2–6 cm wide, base amplexicaul, margin sometimes crenulate, usually discoloured with upper surface green or slightly glaucous and the lower surface copiously white-waxy.

Adult leaves alternate, petiole 1.5–3.8 cm long; blade lanceolate to falcate, 13–35 cm long, 1.5–4.5 cm wide, base tapering to petiole, concolorous, glossy, green, side-veins usually greater than 45° to midrib, densely reticulate, intramarginal vein parallel to and well removed from margin, oil glands island and intersectional.

Inflorescence axillary unbranched, peduncle flattened, 0.3–1.0 cm long, buds 3 per umbel, sessile to stoutly pedicellate (pedicels 0–0.5 cm long). **Mature buds** clavate to pyriform, 0.9–1.2 cm long, 0.7–0.8 cm wide, green to yellow or glaucous, warty, with longitudinal ribs, scar present, operculum flattened and umbonate or beaked, stamens inflexed, anthers cuboid to oblong, versatile, dorsifixed, dehiscing by longitudinal slits (non-confluent), style long, stigma blunt or tapered, locules 3 or 4, the placentae each with 4–6 vertical ovule rows. Flowers white.

Fruit sessile or on pedicels to 0.5 cm long, obconical, 0.6–0.8 cm long, 0.9–1.2(1.6) cm wide, usually slightly angled longitudinally, glaucous or non-glaucous, disc raised and lobed over valves which are about rim level, valves 3 or 4, slightly exerted.

Seeds black, brown or grey, 1.5–3 mm long, ovoid to flattened-ovoid or slightly cuboid, often lacunose, dorsal surface shallowly pitted, hilum ventral.

Cultivated seedlings (measured at ca node 10): cotyledons bilobed; stems square in cross-section and prominently winged also, glaucous; leaves sessile, opposite for many nodes, oblong to ovate or elliptical, 7–10 cm long, 2.5–5 cm wide, amplexicaul, margin entire or subcrenulate, apex pointed, glaucous to blue to blue-grey.


Flowering Time

Flowering has been recorded in February.

Forms of *Eucalyptus globulus* have commonly become naturalised in areas away from the species natural distribution in many places in southern Australia.

Notes

Eucalyptus globulus is a forest tree species that is widespread in the ranges and subcoastal forests of eastern New South Wales as far north as the Carrai Plateau, eastern, southern and central Victoria, and Tasmania. It is notable for the very conspicuous seedlings, coppice



and young saplings with square stems and large, glaucous, oblong to ovate sessile juvenile leaves. The trunks are mostly smooth and the adult leaves are large, bright glossy green and usually falcate.

There are four subspecies:

E. globulus subsp. **globulus**

Of largely lowland distribution in Tasmania, including King Island and islands of the Furneaux group and Rodondo Island, and coastal and subcoastal distribution on the southernmost extremities in Victoria. It has a single, large, sessile, glaucous, warty bud per axil (rarely in threes) and the solitary fruit 1.4–2.7 cm wide.

E. globulus subsp. **bicostata**

A montane and tableland subspecies in eastern New South Wales occurring as far north as the Carrai Plateau, and in Victoria mostly on the slopes of the Great Dividing Range from Orbost in the east west to the Pyrenees, and with a single very old population in South Australia on Mt Bryan north of Burra. It has similar sessile buds to subsp. *globulus*, but they are slightly smaller and occur in threes, and fruit (1)1.3–2.2 cm wide.

E. globulus subsp. **pseudoglobulus**

Mainly in Victoria, from the coastal ranges in east Gippsland but with inland populations at Lerderderg Gorge and north of Toongabbie, and also in Nadgee Nature Reserve, New South Wales. It is morphologically close to subsp. *bicostata* but differs from that subspecies in having narrower fruit: 0.9–1.2(1.6) cm wide in subsp. *pseudoglobulus* and often at least the central bud of the 3-flowered umbel being shortly pedicellate (buds always sessile in subsp. *bicostata*).

E. globulus subsp. **maidenii**

Occurs in subcoastal ranges of far south-eastern New South Wales and far eastern Victoria and has smaller, pedicellate buds in sevens, often not glaucous, but still with the warty, umbonate operculum typical of the group.

Intergradation commonly occurs between subsp. *globulus*, *bicostata* and *pseudoglobulus* where they come into contact, and with populations in southern Victoria it may be impossible to attribute a specimen to any particular subspecies. Also the natural distribution of subsp. *globulus* in Victorian and Tasmanian forests has undoubtedly been confused by forestry plantings in the twentieth century.

Eucalyptus globulus belongs in *Eucalyptus* subgenus *Symphyomyrtus* section *Maidenaria*, a large group of species more or less restricted to south-eastern Australia, characterised by bilobed cotyledons, simple axillary inflorescences, buds with two opercula, stamens with versatile anthers and flattened seeds with a ventral hilum. Within this section, *E. globulus* belongs in series *Globulares* subseries *Euglobulares*, having large, sessile, glaucous, juvenile leaves opposite for many pairs on square, winged stems, buds solitary or in clusters of three or seven and fruit with prominent disc partly covering valves.

Origin of Name

Eucalyptus globulus: Latin *globulus*, a globe, a ball, referring to the fruit shape.

subsp. *pseudoglobulus*: Latin *pseudo-*, false and *globulus*, referring to its resemblance to *E. globulus*.

Copyright © CANBR 2020, all rights reserved.



Web edition hosted at <https://apps.lucidcentral.org/euclid>