

Eucalyptus tereticornis subsp. *basaltica*

Classification

Eucalyptus | Symphyomyrtus | Exsertaria | Erythroxylon

Nomenclature

Eucalyptus tereticornis subsp. **basaltica** A.R.Bean, *Austrobaileya* 9(1): 109-111, fig. 1 (2013).

T: Queensland: Darling Downs District: Near Bottle Tree Lookout, Bunya Mountains National Park, 21 May 1996, P. Grimshaw PG2468 & L.M. Grimshaw; Holotype: BRL; isotype: NSW.

Description

Tree to 30 m tall. Forming a lignotuber.

Bark smooth throughout, pale grey to bluish grey, brownish pink and creamy white mottled; sometimes with persistent dark grey to brown slabs of rough bark on base of older trees.

Juvenile growth (coppice or field seedlings to 50 cm): stem square in cross-section, not winged; juvenile leaves always petiolate (petioles up to ca 1.5 cm long), alternate, ovate to deltoid, 5–11 cm long, 3.5–7.5 cm wide, base usually tapering or rarely truncate, margin entire, apex rounded or bluntly pointed, dull, green to blue-green. (Blade of juvenile leaves held in vertical plane, cf. *E. amplifolia*).

Adult leaves alternate, petiole 1–2.2 cm long; blade lanceolate to falcate, 9–17.5 cm long, 1–3.2 cm wide, base tapering to petiole, margin entire, apex pointed, concolorous, glossy or dull, green, side-veins about 45° to midrib, moderately to densely reticulate, intramarginal vein present, oil glands island.

Inflorescence axillary unbranched, peduncles 0.4–1.4 cm long, more or less terete, buds 7, 9 or 11 per umbel, usually pedicellate (pedicels 0.2–0.6 cm long). **Mature buds** ovoid (0.8–1.2 cm long, 0.4–0.6 cm wide), green to cream, the hypanthium short, operculum bluntly conical (0.6–0.9 cm long and 2 to 3 times length of hypanthium), scar present (outer operculum shed early), stamens erect in bud, anthers cuboid to oblong, versatile, dorsifixed, dehiscing by longitudinal slits (non-confluent), style long, stigma blunt, locules 3, 4 or 5, the placentae each with 6 vertical ovule rows with central gap. Flowers white.

Fruit pedicellate (pedicels 0.2–0.6 cm long), hemispherical, 0.2–0.5 cm long, 0.5–0.8 cm wide, disc steeply ascending obliquely to vertically, valves 3, 4 or 5, strongly exserted.

Seeds black, 1–2 mm long, pyramidal or cuboid, dorsal surface usually pitted, ridges or angles minutely roughened (toothed), hilum terminal.

Cultivated seedlings (measured at ca node 10): cotyledons reniform to oblong; stems squared in cross-section, rarely weakly winged; leaves becoming alternate by nodes 5 to 7, always petiolate, ovate, 6–9.5 cm long, 3.5–5 cm wide, base tapering or rounded, margin entire, apex bluntly pointed, dull, green.

Flowering Time

Flowering has been recorded in May, June, July, August, September, October and November.

Notes

Eucalyptus tereticornis is a species of forest tree widespread in eastern Australia and possibly southern New Guinea. It is characterised by its smooth bark, buds in clusters of 7 to 11 in the leaf axils, stamens held erect inside the buds, fruit with steeply raised disc and exserted valves, typically black, single-coated seed with roughened (toothed) margins; and dull broadly ovate juvenile leaves held in the vertical plane (not horizontal as in *E. amplifolia*).

In the classification of Brooker (2000) *Eucalyptus tereticornis* belongs in the group of red gums (series *Erythroxylon*) which is distinguished by having buds with the stamens mostly erect, fruit where the disc is united to the ovary roof, valves conspicuously exserted and by the black, toothed, cuboid to pyramidal single-coated seed. There are 15 species belonging to this group: *E. amplifolia*, *E. blakelyi*, *E. chloroclada*, *E. dealbata*, *E. dwyeri*, *E. flindersii*, *E. gilleni*, *E. glaucina*, *E. infera*, *E. kabiana*, *E. nandewarica*, *E. nudicaulis*, *E. tereticornis*, *E. terrica* and *E. vicina*.

Eucalyptus tereticornis has been divided into 4 subspecies based on bud shape and juvenile growth.

E. tereticornis subsp. **tereticornis**

Has a horn-shaped operculum dilated at its base; operculum 4–7 times length of base (hypanthium).

A medium-sized to tall forest tree widespread in eastern Australia from Bega on the South Coast of New South Wales to Cape York



Peninsula in north Queensland. Note that plants with buds matching this subspecies do occur in Gippsland, Victoria occasionally, but usually have seed either with two seedcoats like *E. camaldulensis*, or have pied seedcoats (black seedcoat overlaid with a partially developed yellow second seedcoat) suggesting plants of intermediate morphology.

E. tereticornis* subsp. *basaltica

Has ovoid buds with conical operculum not dilated basally and only 2–3 times the length of the hypanthium.

Occurs from south-east Queensland to the Sydney area of New South Wales frequently on basaltic substrates; trees with similar buds occur sporadically further south to Cobargo but are much less common than subsp. *tereticornis*. Seed always black and new juvenile growth green, never glaucous.

E. tereticornis* subsp. *mediana

Usually has much shorter, fatter buds than subsp. *tereticornis* with a broadly conical operculum that isn't basally dilated (operculum 2–4 times length of hypanthium).

Endemic to Gippsland in eastern Victoria, occurring from the Tambo River valley west to Sale and Licola. It grows near *E. camaldulensis* subsp. *camaldulensis* in part of its distribution, e.g. from Bairnsdale to Longford and Rosedale and it is possible that subsp. *mediana* originated from hybrids between *E. tereticornis* and *E. camaldulensis*. Seedlings of subsp. *mediana* have ovate leaves like typical subsp. *tereticornis* and show little variation while those of *E. camaldulensis* from the Sale area and elsewhere are typically lanceolate and vary little. The seeds of subsp. *mediana* can show characters of both *E. tereticornis* (black single-coated seed) and *E. camaldulensis* (yellow double-coated seed) and sometimes seed with only a partially developed second seedcoat thus appearing pied. Juvenile growth of subsp. *mediana* is often glaucous but this is soon lost as leaves mature. The populations of subsp. *mediana* were once widespread on the Gippsland Plain but are now reduced to remnants, mostly in poor health with sparse crowns and few buds and fruit. The best stands are at Briagolong and Moormurung. Similar obese-budded plants occasionally occurring on the South Coast of New South Wales, e.g. at Meringo, Tathra and near Guerilla Bay, have juvenile leaves similar to *E. amplifolia* and are possibly of intermediate morphology.

E. tereticornis* subsp. *rotunda

The buds have operculum not dilated basally, usually with rounded blunt tip and 2–3 times the length of the hypanthium.

Restricted to central Queensland from areas of the Carnarvon Range above 500 m altitude, and possibly the Expedition Range also. The juvenile growth of this subspecies is very different: stems square in cross-section and strongly winged also, leaves broadly ovate to orbicular, prominently petiolate, green, held in vertical plane. This subspecies may be worth species rank.

Within series *Erythroxylon*, trees of *E. tereticornis* are most readily confused with *E. amplifolia* and *E. blakelyi*. *E. amplifolia* has distinctive deltoid to orbicular glossy green juvenile leaves, held horizontally into the juvenile and sapling stage, whilst the juveniles of *E. tereticornis* (and *E. blakelyi*) are broadly ovate and dull and are held in the vertical plane. Also, *E. amplifolia* is a smaller tree and its buds have an operculum lacking the often swollen base compared with typical *E. tereticornis* subsp. *tereticornis*. *E. blakelyi* has a horn-shaped operculum similar to that of *E. tereticornis* subsp. *tereticornis*, and the juvenile leaves of both these species are indistinguishable. *E. blakelyi* is a woodland tree with a more inland distribution than *E. tereticornis* although the two may be difficult to distinguish in far northern New South Wales in the Yetman area, where populations of both are thought to overlap – there is no easy solution.

Two other Queensland red gum species have buds similar to typical *E. tereticornis*: *E. kabiana*, from Mt Beerwah in the Glasshouse Mountains, differs in its mallee habit and narrower lanceolate juvenile leaves. *E. infera* from the Inglewood to Warwick area is also a mallee but has small glossy dark green adult leaves and rounded and emarginate glossy green juvenile leaves.

Of the remaining 10 species in series *Erythroxylon*, all have buds where the conical to rounded operculum is never greater than twice the length of the base of the bud (hypanthium) thus giving an ovoid or diamond-shaped appearance. Six species are woodland trees or mallees of drier tableland or inland areas of New South Wales and/or Queensland: *E. dealbata*, *E. chloroclada*, *E. vicina*, *E. dwyeri*, *E. nandewarica* and *E. terrica*. One species, *E. glaucina*, is restricted to the wetter Hunter Valley area of New South Wales. The last three species are each of restricted distribution in the rocky ranges of the arid zone: *E. flindersii* in north-eastern South Australia, *E. gillenii* in Central Australia, and *E. nudicaulis* in the Mount Isa area of Queensland.

Within its area of occurrence, *E. tereticornis* may be confused with other smooth-barked red gums from the series *Liberivalvae*, viz. *E. bancroftii*, *E. interstans*, *E. prava*, *E. parramattensis* and *E. seeana*. *E. tereticornis* can be distinguished by its fruit which have a steeply ascending disc that is fused to the ovary roof. Species in series *Liberivalvae* all have fruit with an annular disc that is visibly free from the ovary roof (i.e. there is a gap between the disc and the base of the valves in the dehisced fruit).

MORE ABOUT RED GUMS AND OTHER ASSOCIATED GROUPS

Origin of Name

Eucalyptus tereticornis subsp. *basaltica*: the subspecies name refers to the basalt substrate on which these plants commonly grow.



Web edition hosted at <https://apps.lucidcentral.org/euclid>