

# *Eucalyptus vegrandis* subsp. *recondita*

## Classification

Eucalyptus | Symphyomyrtus | Bisectae | Glandulosae | Clinatae

## Nomenclature

**Eucalyptus vegrandis** subsp. **recondita** Nicolle & Brooker, *Nuytsia* 15: 423 (2005).



T: Western Australia: 2.3 km along N boundary fire trail from Bluff Knoll Rd, Stirling Range National Park, 7 Oct. 1982, *Brooker 7695*; holotype: CANB; isotype: PERTH

## Description

**Mallee** to 6 m tall. Forming a lignotuber.

**Bark** smooth throughout, shedding in short ribbons, grey to green-bronze over pink or orange.

**Branchlets** have oil glands in the pith.

**Juvenile growth (coppice or field seedlings to 50 cm):** not seen.

**Adult leaves** alternate, petioles 0.3–1.5 cm long; blade elliptic to oblanceolate, 3.7–6(7.8) cm long, (0.8)1.3–2.5 cm wide, base tapering to petiole, margin distally notched or entire, apex pointed or rounded, glossy, green, surface may appear "glazed" surface, side-veins and reticulation obscured by numerous round oil glands, intramarginal vein present or obscure.

**Inflorescence** axillary unbranched, peduncles erect to spreading, usually flattened, 0.6–1.7 cm long, buds 7, pedicellate (pedicels 0.1–0.5 cm long). **Mature buds** elongated with hypanthium usually just wider than the operculum at the join, 0.7–1.2 cm long, 0.3–0.5 cm wide, scar present (outer operculum lost very early), inner operculum not usually warty, horn-shaped to bluntly conical, hypanthium with weak ridges extending from pedicel, outer stamens erect or terminally deflexed, inner stamens inflexed, anthers oblong, versatile, dorsifixed, dehiscent by longitudinal slits, style long and straight, stigma blunt, locules 4(5), the placentae each with 4 vertical rows of ovules. Flowers white to creamy white.

**Fruit** pedicellate (pedicels 0.1–0.6 cm long), cupular to obconical, 0.6–0.8(1.2) cm long, 0.5–0.8 cm wide, disc descending, valves 4(5), at rim level or slightly exserted.

**Seeds** brown, 1–1.5 mm long, flattened-ovoid, dorsal surface clearly but shallowly reticulate, hilum ventral.

**Cultivated seedlings (measured at node 10):** cotyledons Y-shaped (bisected); stems rounded in cross-section, smooth or slightly warty; leaves shortly petiolate, opposite for 2 to 5 nodes then alternate, linear to lanceolate, 5.5–7 cm long, 0.4–1.6 cm wide, dull grey-green to green for ca 10 nodes then becoming glossy.

## Flowering Time

Flowering has been recorded in early spring, June and February.

## Notes

*Eucalyptus vegrandis* is a mallee species endemic to Western Australia from low-lying sites, often around flats or depressions dominated by saltbush, from Jerramungup west to Katanning and south to Cranbrook and Boxwood Hill. The stems are smooth and the adult leaves glossy, green appearing glazed. Typically it has at least the inner stamen inflexed in bud and the leaf venation obscured by numerous oil glands. Leaf width varies across the range with the most southerly and westerly populations having the widest leaves.

Nicolle & Brooker (2005) described broad-leaved mallees in the south and south-west of the range of *E. vegrandis* as a distinct subspecies, viz. *Eucalyptus vegrandis* subsp. *recondita*. The subspecies can be distinguished as follows:

### ***E. vegrandis* subsp. *vegrandis***

Adult leaves linear to narrowly elliptical, (0.3)0.4–1(1.3) cm wide; with a length:width ratio in the range 6.4–14.2. Occurs from Ongerup to Katanning, Nyabing and Jerramungup. In the field subsp. *vegrandis* is most likely to be confused with, and co-occurs with, the more widespread *E. orthostemon* which has fully erect stamens in bud.

### ***E. vegrandis* subsp. *recondita***

Adult leaves broad, elliptical to oblanceolate, (0.8)1.3–2.5 cm wide, with a length:width ratio in the range 2.1–7.9. Occurs from Cranbrook eastwards around the fringes of the Stirling Range, but not on the Stirling Range itself. Most likely to be confused with broader-leaved forms of *E. suggrandis* subsp. *suggrandis*, and with *E. thamnoides*, both of which have fully erect stamens in the bud.

*Eucalyptus vegrandis* belongs in *Eucalyptus* subgenus *Symphyomyrtus* section *Bisectae* subsection *Glandulosae* because the buds have an operculum scar, cotyledons are bisected and branchlets have oil glands in the pith. Subsection *Glandulosae* is a very large group of ca

80 species. *E. vegrandis* occupies an uncertain position as it appears to be related to species in two different sub-groups in subsection *Glandulosae*: one group being the mallet *E. spathulata* and the mallee *E. orthostemon*, both with very narrow leaves, small similarly shaped buds cf. *E. vegrandis* but with erect stamens inside the bud, and occupying a similar saline habitat, plus *E. suggrandis* subsp. *suggrandis*, a mallee species from heathlands with erect stamens in slightly larger, warty buds; and the other group comprising the mallet *E. cernua* and mallee *E. proxima*, both of which have completely inflexed stamens in bud but differ from *E. vegrandis* in wider leaves and larger, 2-ribbed buds and fruit and colourful flowers.

Correct identification of *E. vegrandis* requires inspecting the arrangement of stamens inside the bud by sectioning it longitudinally and looking at it with a hand lens—the outer stamens will be more or less erect but the inner stamens will be inflexed. The plant habit, especially the presence or absence of a lignotuber, will also need to be known accurately.

Plants of *Eucalyptus vegrandis* in the area from Beaufort Inlet to the Bremer River may be intermediate between the subspecies.

### Origin of Name

*Eucalyptus vegrandis*: Latin *vegrandis*, not very large, referring to both its small stature and because it has been confused with *E. suggrandis*.

Subsp. *recondita*: Latin *reconditis*, meaning hidden or concealed, referring to the fact of its confusion with other species in the group of species related to *E. spathulata*.

---

Copyright © CANBR 2020, all rights reserved.



Web edition hosted at <https://apps.lucidcentral.org/euclid>