

WATTLE

Acacias of Australia

Acacia adunca A.Cunn. ex G.Don



Source: Australian Plant Image Index (a.3428).
ANBG © M. Fagg, 1978



Source: Australian Plant Image Index (a.3431).
ANBG © M. Fagg, 1978



Source: Australian Plant Image Index (a.19289).
ANBG © M. Fagg, 1978



Source: Australian Plant Image Index (a.19290).
ANBG © M. Fagg, 1978



Source: Australian Plant Image Index (a.30922).
ANBG © M. Fagg, 2000



Source: Australian Plant Image Index (dig.21313).
ANBG © M. Fagg, 2011



Source: Australian Plant Image Index (dig.22011).
ANBG © M. Fagg, 2011



Source: Australian Plant Image Index (dig.22012).
ANBG © M. Fagg, 2011



Source: Australian Plant Image Index (dig.24977).
ANBG © M. Fagg, 2011



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com



Acacia adunca occurrence map.
Occurrence map generated via Atlas of Living Australia (<https://www.ala.org.au>).

Common Name

Wallangarra Wattle, Cascade Wattle

Family

Fabaceae

Distribution

Geographically restricted along the Great Divide from Bolivia Ra. N to Legume in north-eastern N.S.W., W to Amiens State Forest (36 km NW of Stanthorpe), in south-eastern Qld.

Description

Bushy **shrub** or **tree** to 6 m high. **Branchlets** slender, dark reddish, **glabrous**. Phyllodes narrowly **linear**, 7–14 cm long, 1–2.5 mm wide, **uncinate** to subuncinate, not **rigid**, commonly finely wrinkled when dry, **glabrous**; **midrib** subimpressed; **lateral nerves** obscure or absent; **gland** 1 or occasionally 2, not overly prominent, slightly **exserted**, the lowermost 1–4.5 cm above **pulvinus**. Inflorescences 4–13-headed racemes; **raceme axes** 15–40 mm long, **glabrous**; heads showy, **globular**, subdensely 9–14-flowered, bright golden; bracteoles \pm **peltate**, with laminae subcircular and golden- or white-**fimbriolate**; peduncles 3–6 mm long, **glabrous**. Flowers 5-merous; sepals united. Pods **rounded** on opposite sides over alternate seeds, to 13 (-18) cm long, 10–12 mm wide, thinly **coriaceous**, **glabrous**. Seeds **longitudinal**, **oblong** to **elliptic**, 5.5–6 mm long, somewhat shiny, blackish, arillate; **funicle** short.

Habitat

Grows in elevated country, in woodland or forest, in sand or sandy loam over granite.

Specimens

Qld: Amiens, *M.Bengston 5134* (BRI); Bald Rock Ck, N of Wallangarra, *E.J.Carroll & I.R.Telford 707* (CANB). N.S.W.: Shiny Ra., Bolivia, *E.F.Constable 7089* (NSW, PERTH).

Notes

Closely allied to *A. betchei* and *A. linearifolia*. Also resembling *A. acrionastes*, *A. floydii* and *A. macnuttiana*. *Acacia adunca* putatively hybridizes *A. neriifolia* in the Girraween NP, southeast Qld.

The complex nomenclatural history of the names *A. adunca* and *A. linearifolia* is discussed by B.R.Maslin, *Telopea* 6: 43–49 (1994). G.Bentham, *Fl. Austral.* 2: 372 (1864), erroneously treated *A. adunca* as conspecific with *A. crassiuscula*. The nomenclature of *A. hamiltoniana* is linked to that of *A. adunca* and is discussed by L.Pedley, *Austrobaileya* 1: 286 (1980) and B.R.Maslin, *Telopea* 6: 43–49 (1994).

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

Minor edits by B.R.Maslin & J.Rogers

B.R.Maslin

This identification key and fact sheets are available as a mobile application:



Australian Government
Department of the Environment and Energy



Department of
Biodiversity, Conservation
and Attractions
Western Australian Herbarium



Australian
Biological
Resources
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.