

# WATTLE

## *Acacias of Australia*

### *Acacia ancistrophylla* C.R.P.Andrews var. *ancistrophylla*



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Acacia ancistrophylla var. *ancistrophylla*  
occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.ala.org.au>).

#### Family

Fabaceae

#### Distribution

Scattered from Wubin (c. 22 km N of Dalwallinu) S to Newdegate (c. 50 km E of Lake Grace), E to Cundeelee Mission (c. 200 km E of Kalgoorlie), and near Salmon Gums, southern W.A.

#### Description

Phyllodes **ascending**, **ob lanceolate to oblong-ob lanceolate, straight**, (1–) 1.2–2.5 (–4) cm long, (1.5–) 2.5–4 mm wide, l:w = 3–8 (–15), with **acute curved to uncinate apex**; nerves obscure; **stomata** obscure. Peduncles 2–5 (–6) mm long; heads 15–23-flowered; sepals free. Pods 1.5–3 mm wide, **straight to shallowly curved, puberulous**. Seeds **oblong**, dull tan; **aril** apical, conical, white.

#### Habitat

Grows in woodland and mallee communities on flats, hillsides and ridges in loam, clay or sandy clay.

#### Specimens

W.A.: 10 km E of Kulja towards Mollerin, *B.R.Maslin 4446* (PERTH); near Coolgardie, Sept. 1920, *C.A.Gardner s.n.* (PERTH); 12.7 km W of Newdegate, *J.W.Green 4464* (PERTH); Cowcowing, *M.Koch 1034* (PERTH);

#### Notes

One collection (Pingaring, *N.Perry 548*, PERTH) has ±pungent phylode tips, similar to those of *A. amyctica*, but the nerves are obscure as in var. *ancistrophylla*. The J.W.Green collection cited above has strongly curved phyllodes, some even sigmoidally curved, and a higher proportion of tetramerous flowers than is typical. The Koch collection cited has glabrous peduncles that are longer than usual.

Nearest variety is var. *perarcuata* which has ±coiled pods; also closely related to *A. amyctica* which has sharply to coarsely pungent phyllodes.

#### FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

#### Author

R.S.Cowan, B.R.Maslin

Minor edits by B.R.Maslin & J.Reid

---

This identification key and fact sheets are available as a mobile application:



Australian Government  
Department of the Environment and Energy



Department of  
Biodiversity, Conservation  
and Attractions  
Western Australian Herbarium



Australian  
Biological  
Resources  
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>  
Copyright 2018. All rights reserved.