

# WATTLE

## *Acacias of Australia*

### *Acacia aspera* Lindl. subsp. *aspera*



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin




Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin




Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: Australian Plant Image Index (a.9920).  
ANBG © M. Fagg, 1991



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Acacia aspera subsp. aspera occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.ala.org.au>).

## Family

Fabaceae

## Distribution

Occurs from The Grampians to Warby Ra., Vic, northern A.C.T. and from Peak Hill S to the Ardlethan-Temora area, and at Sutton Forest, N.S.W.

## Description

Phyllodes 10–30 mm long, flat to slightly concavo-convex in section. Peduncles (1–) 2–10 mm long, stout, mostly with sessile or subsessile glands; heads 30–50-flowered, 6–9 mm diam., bright golden yellow. Bracteoles eglandular, exserted in the buds (buds usually burr-like in appearance).

## Specimens

N.S.W.: Jindalee, Sutton Forest, *W. de Beuzeville* (NSW252801 *n.v.*); Herveys Ra. (sphalm. 'Barker Ra.'). between Baldy and Peak Hill, *E. Gordon* (NSW78949 *n.v.*). A.C.T.: Ginninderra, 26 Aug. 1968, *C. Burgess* (CANB *n.v.*). Vic.: Bendigo Whipstick, c. 12 km N of Bendigo, *B.R. Maslin 5849* (PERTH); between Benalla and Glenrowan, 19 km SW of Wangaratta, *N. Hall H79/98* (NSW, PERTH).

## Notes

Subsp. *aspera* is not sympatric with subsp. *parviceps* but the two occur as close as c. 40 km apart in a few localities.

## FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

## Author

Revised by B.R.Maslin

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>  
Copyright 2018. All rights reserved.