

# WATTLE

## Acacias of Australia


### *Acacia basedowii* Maiden



Image courtesy of Northern Territory Herbarium



Image courtesy of Northern Territory Herbarium



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Acacia basedowii occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.ala.org.au>).

### Common Name

Basedow's Wattle

### Family

Fabaceae

### Distribution

Found in central Australia from Rawlinson Ra. and Skirmish Hill, W.A., E to Macdonnell Ra., N.T. and Musgrave Ra., S.A.

### Description

Very similar to the more westerly distributed *A. oxyclada* but distinguished by branchlets with a conspicuous, flaking, white epidermis, phyllodes 5–20 mm long and 1–2 mm wide, peduncles 5–12 mm long and glabrous to subglabrous, and heads commonly 25–35-flowered; only old pod valves and no seeds seen.

### Habitat

Occurs on rocky slopes of ranges.

### Specimens

W.A.: Rawlinson Ra., J.R.Maconochie 815 (PERTH). N.T.: Mt Fraser, Mulga Park Stn, T.S.Henshall 167 (NSW, PERTH). S.A.: slopes of mountain E of Mt Woodroffe, Musgrave Ra., D.E.Symon 2675 (AD, K).

### Notes

One of the diaphyllodinous species (see discussion under *A. diaphyllodinea* for definition and references); probably better treated as an infraspecific taxon of *A. oxyclada*, but seeds should be examined before any change is made.

### FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

### Author

B.R.Maslin

Minor edits by J.Rogers

This identification key and fact sheets are available as a mobile application:

