


# WATTLE

## *Acacias of Australia*

### *Acacia baueri subsp. *aspera* (Maiden & Betche) Pedley*



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Acacia baueri subsp. *aspera* occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.ala.org.au>).

#### Family

Fabaceae

#### Distribution

Uncommon in N.S.W. Occurs S of Sydney, Royal Natl Park and near Wilton, and at Wentworth Falls in the Blue Mtns.

#### Description

Branchlets puberulous, tuberculate. Phyllodes in irregular whorls or scattered, rather crowded, with scattered tubercle-based hairs when young. Peduncles 2–4 mm long, usually hidden by the phyllodes. Heads normally 8–10-flowered.

#### Habitat

Occurs in coastal low heath, grows in rocky sandstone areas.

#### Specimens

N.S.W.: Royal Natl Park on the road to Warrumbul, R.Coveny 4211 et al. (NSW); 'Little Switzerland', Kings Tableland, T.Tame 1925 (PERTH).

#### Notes

The low stature, ebracteate peduncle bases and striate pods of *A. baueri* subsp. *aspera* are reminiscent of *A. bynoeana* which differs markedly in its flat, 3-nerved, pungent phyllodes. Sometimes confused with *A. gordoni*.

#### FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

#### Author

B.R.Maslin

Minor edits by B.R.Maslin

This identification key and fact sheets are available as a mobile application:



Australian Government  
Department of the Environment and Energy



Department of  
Biodiversity, Conservation  
and Attractions  
Western Australian Herbarium



Australian  
Biological  
Resources  
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>  
Copyright 2018. All rights reserved.