


WATTLE

Acacias of Australia

Acacia baueri Benth. subsp. *baueri*



Source: Australian Plant Image Index (dig.3153).
ANBG © M. Fagg, 2006



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com



Acacia baueri subsp. *baueri* occurrence map.
Occurrence map generated via Atlas of Living
Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Occurs from Fraser Is., Qld, S to Sydney, N.S.W. The subspecies is uncommon and is in danger of extinction in the extreme south-east of Qld and in the Sydney area, *vide* L.Pedley, *Contr. Queensland Herb.* 11: 10 (1972).

Description

Branchlets **glabrous** or **puberulous**, sometimes **tuberculate**. Phyllodes in regular whorls 0.5–2 cm apart, smooth or occasionally **tuberculate**. Peduncles 5–16 mm long. Heads 10–20-flowered.

Habitat

Found in low-lying coastal heath on infertile and often waterlogged sands.

Specimens

Qld: Hollywell near Southport, *L.Pedley 2178* (NSW). N.S.W.: Chinaman's Beach, 3.2 km SE of Evans Head, *R.Coveny 4592* (PERTH); near Coffs Harbour, *T.Tame 1716* (PERTH).

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

B.R.Maslin

Minor edits by B.R.Maslin

This identification key and fact sheets are available as a mobile application:



Australian Government
Department of the Environment and Energy



Department of
Biodiversity, Conservation
and Attractions
Western Australian Herbarium



Australian
Biological
Resources
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.