

# WATTLE

## Acacias of Australia

### *Acacia beauverdiana* Ewart & Sharman



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
J. & M. Simmons




Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Acacia beauverdiana occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.ala.org.au>).

### Family

Fabaceae

### Distribution

Occurs from near Bonnie Rock S to near Lake Grace with a disjunct occurrence near Broad Arrow, south-western W.A.

### Description

Shrub 1–4 m high, rounded, occasionally a tree 3–8 m high. Branchlets terete, glabrous, sometimes hairy in phyllode axils. Phyllodes ascending to erect, linear to narrowly oblong-elliptic, ±straight, rarely terete, 7–13 cm long, usually 1.5–3 mm wide, with a short-acuminate and gently curved to uncinata apex, glabrous except pulvinus sometimes hairy adaxially, with numerous very fine closely parallel nerves; central nerve sometimes slightly more evident than the rest. Inflorescences simple, mostly 2 per axil; peduncles 1.5–4 mm long, ±puberulous; heads short-obloid to globular, 6–8 mm long, 5–7 mm diam., 28–36-flowered, golden. Flowers 5-merous; sepals 2/3-united. Pods linear, raised over and constricted between seeds, straight or slightly curved, to 9 cm long, 2.5–3 mm wide, chartaceous, glabrous. Seeds longitudinal, narrowly oblong-elliptic, 2.5–4 mm long, glossy, tan; funicle/aril fleshy, folded several times to form long conical terminal aril.

### Habitat

Grows in sand, sandy gravel and loam, in heath, scrub, shrubland and low open woodland.

### Specimens

W.A.: 12.5 km W of Moorine Rock on Great Eastern Hwy, B.R.Maslin 4477 (MEL, PERTH); 25 km NE of Bonnie Rock, K.Newbey 9621 (PERTH); 2.7 km W of Merredin, M.D.Tindale 3729 (NSW, PERTH).

### Notes

K.Newbey 9621 from Bonnie Rock at the northern limit of the geographic range is atypical in having phyllodes 5–6 mm wide. R.Cumming 2457 (CANB, PERTH) from Yorkrakine is atypical in having terete phyllodes.

### FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

### Author

R.S.Cowan, B.R.Maslin

This identification key and fact sheets are available as a mobile application:



Australian Government  
Department of the Environment and Energy



Department of  
Biodiversity, Conservation  
and Attractions  
Western Australian Herbarium



Australian  
Biological  
Resources  
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>  
Copyright 2018. All rights reserved.