

WATTLE

Acacias of Australia

Acacia beckleri Tindale subsp. *beckleri*



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com



Acacia beckleri subsp. *beckleri* occurrence map.
Occurrence map generated via Atlas of Living
Australia (<https://www.ala.org.au>).

Common Name

Barrier Range Wattle

Family

Fabaceae

Distribution

Occurs from the Gawler Ra. and NE Eyre Penin, S.A., E to the Barrier Ra. and the Manara Hills–Ivanhoe area, N.S.W.; also further E in N.S.W. from near Dubbo, Cobar and Jerilderie but probably naturalized or cultivated in these areas.

Description

Peduncles (2)–4–7 (–8) mm long, 1–2 mm wide, very thick (dry); heads 8–10 mm diam. (dry), (50)–54–67-flowered.

Habitat

Grows in shrubland on rocky ridges and hillsides or sometimes in sand in mulga communities.

Specimens

S.A.: Corunna Hill, Iron Knob, B.Copley 2320 (AD, PERTH); Gawler Ra., c. 5 km from intersection of Yardea, Nonning and Kingoonya Rds, A.E.Orchard 2335 (AD). N.S.W.: Mootwingee, NE of Broken Hill, B.G.Briggs & J.De Nardi B2761 (NSW, PERTH); Jerilderie, late June 1964, D.Waterhouse (NSW).

FOA Reference

Flora of Australia Project

Author

B.R.Maslin

This identification key and fact sheets are available as a mobile application:



Australian Government
Department of the Environment and Energy



Department of
Biodiversity, Conservation
and Attractions
Western Australian Herbarium



Australian
Biological
Resources
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>

Copyright 2018. All rights reserved.