

# WATTLE

## Acacias of Australia

### Acacia blakei subsp. *diphylla* (Tindale) Pedley



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)




Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)



Acacia blakei subsp. *diphylla* occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.ala.org.au>).

## Family

Fabaceae

## Distribution

Occurs in N.S.W. near Wollomombi Falls and Hillgrove on the Northern Tablelands, NNW of Gloucester on the North Coast, and extending southwards to the Coxs R. and Lake Yarrunga; also in south-eastern Qld S of the Ipswich area.

## Description

Phyllodes **elliptic** when immature (not occurring in subsp. *blakei*), and narrowly to very narrowly **elliptic** when mature. Spikes rarely longer than 4 cm. Bracteoles without red-brown **papillae** on the **lamina** or if present only sparsely so. Flowers: **calyx** (mainly tube) ±densely covered in white-**hyaline** hairs, rarely with sparse red-brown **papillae** on apices; **ovary** distinctively **pubescent** at apex and base, usually **glabrous** in middle.

## Phenology

Flowers Aug.–Nov.

## Habitat

Grows usually in shale- or slate-derived soils, common in woodland fringing gorges and on hillsides, often in association with *Eucalyptus tereticornis* and *E. melliodora*.

## Specimens

Qld: Mt Maroon, SW of Rathdowney, A.R.Bean 6669 (BRI n.v., CANB n.v., NSW); Amiens, Sow and Pigs, L.Pedley 4952 (BRI n.v., CANB n.v., DNA n.v., K n.v., MEL n.v., MO n.v., NSW, PERTH n.v., PR n.v.). N.S.W.: Bakers Ck Falls, R.G.Coveny 3927 & K.Thurtell (A, AD, BRI, CANB, K, L, LE, MEL, NSW, PERTH, RSA, U, UC, USA); 22.3 km NNW of Gloucester on Giro Rd, K.Thurtell & R.G.Coveny 3810 (AD, BRI, CANB, CHR, K, L, MEL, NSW, PERTH, US); edge of Hillgrove Gorge, c. 17 miles [27.2 km] SE of Armidale, Oct. 1964, J.B.Williams (NSW).

## FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

## Author

Dr M.D.Tindale and Dr P.G.Kodela with the assistance of M.Bedward, S.J.Davies, C.Herscovitch, D.A.Keith and/or D.A.Morrison

This identification key and fact sheets are available as a mobile application:

