

WATTLE

Acacias of Australia

Acacia brachyphylla var. *recurvata* R.S.Cowan & Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com



Acacia brachyphylla var. *recurvata* occurrence map.
Occurrence map generated via Atlas of Living Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Occurs from near Kukerin SE to near Jerramungup, south-western W.A.

Description

Phyllodes 8–10 mm long; apex **recurved** to subuncinate, strongly **mucronate**; **gland** near **distal** end of **pulvinus** or absent. Pods to 2 cm long, 1.5–3 mm wide, **pruinose**, **glabrous** or subglabrous and **glabrescent**.

Habitat

Grows in often gravelly loam and sand, in shrubland.

Specimens

W.A.: 4 km N of Calyerup Rock, *K.Newbey* 4231 (PERTH); 1.6 km E of Nyabing, *K.Newbey s.n.* (PERTH).

Notes

One collection (*K.J.Atkins* 89008, PERTH) is unusual in having glabrous phyllodes and branchlets.

Superficially similar to *A. vassalii*.

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

R.S.Cowan

This identification key and fact sheets are available as a mobile application:



Australian Government
Department of the Environment and Energy



Department of
Biodiversity, Conservation
and Attractions
Western Australian Herbarium



Australian
Biological
Resources
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.