

WATTLE

Acacias of Australia

Acacia chapmanii subsp. *australis* R.S.Cowan & Maslin



Source: WorldWideWattle ver. 2.

Published at: www.worldwidewattle.com



Acacia chapmanii subsp. *australis* occurrence map.

Occurrence map generated via Atlas of Living Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Known only from near Bolgart, south-western W.A.

Description

Stipules ±1 mm long, not at all rigid, subulate; gland 4–9 mm above phyllode base. Phyllodes ascending, subterete to flat, shallowly recurved. Heads 24–27-flowered; bracteoles evident in buds, with laminae much broader than sepals. Pods and seeds not seen.

Habitat

Grows in sand and sandy gravel in low heath.

Specimens

W.A.: between Bolgart and Calingiri, S.Paust 1005 (PERTH).

FOA Reference


Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

R.S.Cowan

Minor edits by B.R.Maslin & J.Reid

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.