

WATTLE

Acacias of Australia

Acacia dissona R.S.Cowan & Maslin var. *dissona*



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Acacia dissona var. dissona occurrence map.
Occurrence map generated via Atlas of Living
Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Disjunct, occurring from Merredin–Southern Cross S to near Ongerup with outliers in the Coorow–Wanarra area (c. 250 km due NW of Merredin) and the Norseman area (c. 350 km due NE of Ongerup).

Description

Phyllodes subterete, 2–4 cm long, contracted to ±sharply pungent apex; nerves and internerve spaces often uniform in colour; stomata sometimes tuberculate. Pods distinctly constricted between the seeds, sparingly appressed-puberulous. Seeds 4.5 mm long; aril long-conical.

Habitat

Grows in a variety of soils, often clay, in eucalypt woodland and mallee.

Specimens

W.A.: 21.6 km ENE of Merredin, R.Coveny 8340 & B.Habersley (NSW, PERTH); 5 km SE of Coorow, A.Doley 4 (PERTH); Norseman, J.Goodwin 271 (PERTH); c. 4 km S of Karlgarin on Pederah Rd, B.R.Maslin 6738 (PERTH).

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

R.S.Cowan

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.