

Acacia eriopoda Maiden & Blakely x Acacia tumida F.Muell. ex Benth. var. tumida

Family

Fabaceae

Distribution

Most collections are from the immediate vicinity of Broome although it is also recorded from about 100 km N of this township on the Dampier Peninsula and from W of Erskine Range (S of Derby).

Specimens

W.A.: south side of Gantheaume Point, Broome, B.R. Maslin 4921A (PERTH); 100.5 km N of Broom on the road to Beagle Bay Mission, 20 June 1981, B.R. Maslin 4926 (DNA, K, PERTH); Broome, c. 0.5 km S of golf course, B.R. Maslin 7226 (CANB, DNA, PERTH); c. 61 km W of Erskine Range on Great Northern Highway (S of Derby), B.R. Maslin 7314 (NSW, PERTH); near Port Drive Industrial area on 'Riddell Beach Back-track', Broome, T. Willing 68 (MELU, PERTH).

Notes

Morphologically this putative hybrid combines the phyllodes of *A. tumida* var. *tumida* (although often slightly narrower) and the pods of *A. eriopoda*. Both putative parents are recorded as occurring with the putative hybrid in a number of localities.

FOA Reference

Flora of Australia Project

Author

B.R.Maslin

This identification key and fact sheets are available as a mobile application:













URL: https://keys.lucidcentral.org/keys/v3/wattle Copyright 2018. All rights reserved.