

# WATTLE

## *Acacias of Australia*

### *Acacia extensa* Lindl.



Source: Australian Plant Image Index (a.9969).  
ANBG © M. Fagg, 1985



Source: Australian Plant Image Index  
(dig.22020).  
ANBG © M. Fagg, 2011



Source: Australian Plant Image Index  
(dig.22021).  
ANBG © M. Fagg, 2011




Source: Australian Plant Image Index  
(dig.33892).  
ANBG © M. Fagg, 2013



Source: Australian Plant Image Index (dig.8717).  
ANBG © M. Fagg, 2009



Source: Australian Plant Image Index  
(dig.33893).  
ANBG © M. Fagg, 2013



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.



Acacia extensa occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.alb.org.au>).

### Common Name

Wiry Wattle

### Family

Fabaceae

### Distribution

Widely distributed from near Mogumber S to Albany, south-western W.A.

### Description

Spindly open shrub to 3 m high; glabrous, occasionally semi-weeping. Branchlets angled, often flattened at apices, sometimes narrowly winged, green, with yellowish ribs. Phyllodes often few, distant and resembling the branchlets, frequently filiform, slender, normally  $\pm$ pentagonal in section, rarely flat, 6–24 cm long, 0.8–1.8 mm wide, with somewhat indurate mucro, with 5 prominent yellowish nerves (the adaxial 2 forming a broad upper margin). Inflorescences comprising racemes resembling the branchlets, sometimes simple; peduncles 3–8 mm long; heads globular, 12–24-flowered, light golden. Flowers 5-merous; sepals united to near apex; petals 1-nerved. Pods linear, submoniliform, straight to shallowly curved, to 9 cm long, 3–4 mm wide, firmly chartaceous to thinly coriaceous. Seeds longitudinal, oblong to oblong-elliptic, 4–5 mm long, shiny, dark brown; aril apical.

### Habitat

Widely distributed from near Mogumber S to Albany, south-western W.A.; naturalized in a few places in Vic.

### Specimens

W.A.: 3 km S of Porongurup Ra., R.H.Kuchel 1955 (PERTH); c. 3.5 km below Mundaring Weir, Helena R. Valley, B.R.Maslin 671A (CANB, PERTH); Darling Ra., L.Pritzel 532 (B, BM, E, G, L, LY, M, NY, P, PR, Z). Vic.: Western Grampians, Wartook State Forest, G.W.Carr 0908-267 (MEL).

### Notes

Included in sect. *Pulchelloidea* by J.Vassal, *Bull. Soc. Hist. Nat. Toulouse* 108: 1–127 (1972); this section combined both phyllodinous and bipinnate-leaved species.

Besides the names given in synonymy above, a number of other names, presumably unpublished, appear on specimens of *A. extensa* at various European herbaria, viz. *A. balsamea* (WU), *A. balsamifera* (FI), *A. caetinsa* (MPU) and *A. howardii* (L).

### FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

### Author

Minor edits by B.R.Maslin

B.R.Maslin

---

This identification key and fact sheets are available as a mobile application:



Australian Government  
Department of the Environment and Energy



Department of  
Biodiversity, Conservation  
and Attractions  
Western Australian Herbarium



Australian  
Biological  
Resources  
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>  
Copyright 2018. All rights reserved.