

WATTLE

Acacias of Australia

Acacia isoneura subsp. *nimia* Maslin & A.R.Chapm.



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Acacia isoneura subsp. nimia occurrence map.
Occurrence map generated via Atlas of Living
Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Occurs in the Wubin-Perenjori area.

Description

Phyllodes thick, rigid (0.8–1.2 mm diam.); nerves 0.3 mm wide. Spikes single within phyllode axils. Seeds grey-brown speckled dark brown.

Habitat

Grows in sand in tall shrubland.

Specimens

W.A.: 10.5 km from Perenjori on the Three Springs rd, I.B.Armitage 372 (PERTH); 9 km S of Latham on road to Wubin, B.R.Maslin 6434 (CANB, K, MEL, PERTH).

Notes

Similar to *A. cylindrica* which has 16-nerved phyllodes.

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

A.R.Chapman, B.R.Maslin

Minor edits by B.R.Maslin & J.Reid

This identification key and fact sheets are available as a mobile application:



Australian Government
Department of the Environment and Energy



Department of
Biodiversity, Conservation
and Attractions
Western Australian Herbarium



Australian
Biological
Resources
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.