

Acacia leiocalyx subsp. herveyensis Pedley





Acacia leiocalyx subsp. herveyensis occurrence map.

Occurrence map generated via Atlas of Living Australia (https://www.ala.org.au).

Family

Fabaceae

Distribution

Occurs in eastern Qld from Yeppoon SE to Fraser Is. and S to Gympie; it is sympatric with subsp. leiocalyx in the Gympie–Maryborough area.

Description

Phyllodes 9-16 (-18) cm long, 7-14 (-16) mm wide, 8-16 (-19) times as long as wide.

Phenology

Flowers July-Sept.

Habitat

Grows in sand in beach scrub or in clay soils in eucalypt forest.

Specimens

Qld: Boonooroo, *D.Boyland 1507* (BRI); 20 miles [32 km] SW of Double Island Pt, *S.L.Everist 7856* (BRI); Arthurs Lookout near Round Hill Head, *L.Pedley 2873* (BRI, NSW).

FOA Reference

Data derived from Flora of Australia Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

Dr M.D.Tindale and Dr P.G.Kodela with the assistance of M.Bedward, S.J.Davies, C.Herscovitch, D.A.Keith and/or D.A.Morrison, B.R.Maslin

This identification key and fact sheets are available as a mobile application:













URL: https://keys.lucidcentral.org/keys/v3/wattle Copyright 2018. All rights reserved.