

WATTLE

Acacias of Australia

Acacia leiocalyx (Domin) Pedley subsp. *leiocalyx*



Source: Australian Plant Image Index
(dig.23033).

ANBG © M. Fagg, 2012



Source: Australian Plant Image Index
(dig.23629).

ANBG © M. Fagg, 2012



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com

B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com

B.R. Maslin



Source: Australian Plant Image Index
(dig.28055).

ANBG © M. Fagg, 2013



Source: Australian Plant Image Index
(dig.23034).

ANBG © M. Fagg, 2012



Source: Australian Plant Image Index
(dig.23626).

ANBG © M. Fagg, 2012



Source: Australian Plant Image Index
(dig.23627).

ANBG © M. Fagg, 2012



Source: Australian Plant Image Index
(dig.23628).

ANBG © M. Fagg, 2012



Source: Australian Plant Image Index
(dig.28056).

ANBG © M. Fagg, 2013



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com

B.R. Maslin




Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com

B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com

B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com

See illustration.




Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com

See illustration.



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com

See illustration.



Acacia leiocalyx subsp. *leiocalyx* occurrence map.

Occurrence map generated via Atlas of Living Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Widespread in Qld where it extends from S of Mackay (inland to about 146°E) to north-eastern N.S.W.; also the Sydney and Narrabri districts in N.S.W.

Description

Phyllodes 7–12 (–18) cm long, (10–) 12–25 (–35) mm wide, 4–9 times as long as wide.

Phenology

Flowers mid-Apr.–July in coastal Qld; late Apr.–Aug. in inland Qld; (May–) June–Sept. in N.S.W.

Habitat

Grows in eucalypt forest, woodland and heaths, in sandy or clayey soils.

Specimens

Qld: on Ban Ban Springs–Gayndah road, 72 km by road from Biggenden towards Gayndah, *S.J.Davies 59 & J.M.Powell* (AD, K, MEL, NSW, PERTH); 4 miles [6.4 km] WNW of Chinchilla, *M.D.Tindale 716* (AD, CANB, K, NSW). N.S.W.: 3.7 km S of Corindi, *R.G.Coveny 4261 & J.Armstrong* (A, AD, BRI, CANB, G, L, MEL, NSW, TNS, UC, US); 2.4 km SE of Kempsey, *K.Thurtell & R.G.Coveny 3835* (A, AD, CANB, CHR, K, L, LE, MEL, PERTH, RSA, U, US).

Notes

Subspecies *leiocalyx* differs from subsp. *herveyensis* in having generally broader phyllodes and (at least in coastal Qld areas where subsp. *herveyensis* occurs) flowers earlier in the season.

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

Dr M.D.Tindale and Dr P.G.Kodela with the assistance of M.Bedward, S.J.Davies, C.Herscovitch, D.A.Keith and/or D.A.Morrison, B.R.Maslin

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.