

WATTLE

Acacias of Australia

Acacia nigripilosa subsp. *latifolia* Maslin



Source: WorldWideWattle ver. 2.

Published at: www.worldwidewattle.com

See illustration.



Acacia nigripilosa subsp. latifolia occurrence map.

Occurrence map generated via Atlas of Living Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Occurs between Caron and Maya and NE of Geraldton. Terete phyllode variants of subsp. *nigripilosa* occur also in the general region of Caron and Maya.

Description

Phyllodes (1.5–) 2–3.5 cm long, (4–) 5–8 (–10) mm wide, l:w = 3–6, normally dull and glaucous to subglaucous, rather abruptly narrowed at apex to straight, ±pungent point.

Specimens

W.A.: N of Caron, M.D.Tindale 2777 (CANB n.v., K n.v., L n.v., MEL n.v., NSW, PERTH, US n.v.); road to mallee Stn, N of Wandana Nature Reserve, c. 120 km NE of Geraldton, P.C.Ryan 165 (PERTH).

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

B.R.Maslin

Minor edits by J.Rogers

This identification key and fact sheets are available as a mobile application:



Australian Government
Department of the Environment and Energy



Department of
Biodiversity, Conservation
and Attractions
Western Australian Herbarium



Australian
Biological
Resources
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.