

# WATTLE

## *Acacias of Australia*

### *Acacia plectocarpa* A.Cunn. ex Benth. subsp. *plectocarpa*



Source: Australian Plant Image Index (a.31249).  
ANBG © M. Fagg, 1999



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
J. & M. Simmons



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
J. & M. Simmons



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
J. & M. Simmons



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin




Source: Australian Plant Image Index (a.31250).  
ANBG © M. Fagg, 1999



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
Kym Brennan



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.



*Acacia plectocarpa* subsp. *plectocarpa*  
occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.ala.org.au>).

## Family

Fabaceae

## Distribution

Widespread in north-western W.A. through N.T. to far NW Qld, N of 18°S on the coast and up to 150 km inland.

## Description

Phyllodes **linear** to narrowly **elliptic**, **straight** or **curved**, (8.5–) 9.8–20 (–26) cm long, 4.5–15 mm wide (occasionally several phyllodes atypically wider); nerves with 1–3 more prominent than the rest; **pulvinus** 1.5–4 mm long. Spikes bright yellow, 3–4 mm wide. **Calyx** 0.3–0.5 mm long. Pods 6–12 mm wide. Seeds transverse; **funicle** folded 4 times.

## Phenology

Flowers Mar.–July.

## Habitat

Grows in sandy soil mostly near watercourses, in eucalypt woodland on sandstone or laterite.

## Specimens

W.A.: Denham R., *W.V.Fitzgerald* (NSW183370); 21 miles [33.8 km] E of Kununurra, *I.V.Newman* 626 (NSW); Wyndham, Sept. 1903, *A.E.V.Woodroffe* (NSW). N.T.: 10 km N of Pine Creek, *M.D.Tindale* 6003 & *C.Dunlop* (BRI, CANB, K, MEL, NSW, US); 3.8 km S of Edith R., *M.D.Tindale* 6021 & *C.Dunlop* (CANB, DNA, K, NSW, PERTH, UNSW). Qld: 4 km from Hells Gate on Doomadgee road, *P.I.Forster* PIF21074 and *R.Booth* (BRI).

## Notes

Details of ecology, utilisation, etc. of *A. plectocarpa* subsp. *plectocarpa* are given in J.W.Turnbull (ed.), *Multipurpose Austral. Trees & Shrubs* 182 (1986).

## FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

## Author

Minor edits by B.R.Maslin & J.Rogers

Dr M.D.Tindale and Dr P.G.Kodala with the assistance of M.Bedward, S.J.Davies, C.Herscovitch, D.A.Keith and/or D.A.Morrison

This identification key and fact sheets are available as a mobile application:



Australian Government  
Department of the Environment and Energy



Department of  
Biodiversity, Conservation  
and Attractions  
Western Australian Herbarium



Australian  
Biological  
Resources  
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>  
Copyright 2018. All rights reserved.