

WATTLE

Acacias of Australia

Acacia praetermissa Tindale



Source: Australian Plant Image Index
(dig.13295).
ANBG © M. Fagg, 2007



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
J. & M. Simmons



Source: Australian Plant Image Index
(dig.13298).
ANBG © M. Fagg, 2007



Source: Australian Plant Image Index
(dig.13447).
ANBG © M. Fagg, 2007



Source: Australian Plant Image Index
(dig.13296).
ANBG © M. Fagg, 2007



Source: Australian Plant Image Index
(dig.13297).
ANBG © M. Fagg, 2007



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
Kym Brennan



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
Kym Brennan



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
Kym Brennan



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Acacia praetermissa occurrence map.
Occurrence map generated via Atlas of Living
Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Restricted to the Pine Creek–Emerald Springs region of the N.T., although poorly collected.

Description

Multistemmed **shrub** to 1.8 m high, resprouting from perennial rootstock following fire. Bark smooth, bluish grey, blotched. **Branchlets terete**, purplish brown, often lightly **pruinose**; lenticels visible. Phyllodes spreading, very narrowly **ob lanceolate-elliptic, falcate**, 9–15 (–18) cm long, 3–12 (–18) mm wide, with prominent midnerve and 2–4 subprominent parallel nerves; minor nerves 1–3 per mm, loosely **anastomosing**; **gland** 1, basal, adjacent to **pulvinus**. Spikes 1–2.7 cm long, golden. Flowers loosely arranged, 5-merous. **Calyx** 0.5–1 mm long, dissected to 1/4–1/2, sparsely fimbriate towards apex; **corolla** 1.3–1.8 mm long, dissected to 1/3–1/2, **glabrous**; **ovary** sparsely **pilose**. Pods narrowly **ob lanceolate** to **± linear, straight-sided** or slightly constricted between seeds, **± flat**, 4.5–10.5 cm long, **woody**, obliquely to longitudinally and reticulately nerved, opening elastically from apex; margins thickened. Seeds slightly **oblique, elliptic-oblong** or **elliptic to orbicular**, 5.5–7 mm long, brown; **areole** open; **funicle-aril** conical.

Phenology

Flowers Feb.–Aug.

Habitat

Usually on hillsides in lateritic soil or sandy silt in eucalypt woodland.

Specimens

N.T.: 5 miles [8 km] N [of] Pine Creek, N.G.Eddy 120 (DNA, NSW); c. 0.5 km N of Emerald Springs Roadhouse, B.R.Maslin 7348, M.McDonald & G.Leach (NSW, PERTH); 1.5 km N of Emerald Springs, Stuart Hwy, M.D.Tindale 10077, P.Munns & R.T.Turley (B, DNA, MEL, NSW, PERTH).

Notes

Acacia praetermissa can be distinguished from members of the '*A. stigmatophylla* group' by the very loosely anastomosing minor nerves between the 3–5 parallel main nerves, the ridges of the branchlets which are not minutely crenulated, and the compressed-orbicular seeds.

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

Dr M.D.Tindale and Dr P.G.Kodela with the assistance of M.Bedward, S.J.Davies, C.Herscovitch, D.A.Keith and/or D.A.Morrison

Minor edits by B.R.Maslin

This identification key and fact sheets are available as a mobile application:



Australian Government
Department of the Environment and Energy



Department of
Biodiversity, Conservation
and Attractions
Western Australian Herbarium



Australian
Biological
Resources
Study



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.