

WATTLE

Acacias of Australia

Acacia ramulosa var. linophylla (W.Fitzg.) Pedley



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Acacia ramulosa var. linophylla occurrence map.
Occurrence map generated via Atlas of Living
Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Extending from the extreme west of W.A., where it is apparently common, especially in the lower part of the Gascoyne R. basin to S.A., where it occurs sporadically over a wide area.

Description

Phyllodes **terete**, 7.5–13.5 (~16) cm long, 0.7–1.4 mm diam. Spikes 10–30 mm long. **Corolla** 1.5–1.7 mm long, with long **hyaline** hairs usually confined to midlines of lobes. Seeds to 7 mm long and 5 mm wide.

Habitat

It occurs on sandy soils, often on dunes and on sand-plains, though habitat data are lacking for many of its western occurrences. Associated taxa are reported to be *A. ramulosa* var. *ramulosa* and *A. aneura*.

Specimens

W.A.: Yalary Stn, 95 km E of N.W.C.H. [North West Coastal Hwy], B.R.Maslin 5465 (PERTH); 3 miles [5 km] SW of Millrose, N.H.Speck 1377 (BRI, CANB). S.A.: Great Victoria Desert, Serpentine Lakes, 170 km W of Vokes Hill Corner, H.Hewitson (AD, BRI); Barton, C.E.Hubbard 8337 (BRI, K).

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

L.Pedley

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.