

WATTLE

Acacias of Australia

Acacia rigens A.Cunn. ex G.Don



Source: Australian Plant Image Index (a.31304).
ANBG © M. Fagg, 2002



Source: Australian Plant Image Index (dig.15579).
ANBG © M. Fagg, 2009



Image courtesy of Northern Territory Herbarium



Source: Australian Plant Image Index (dig.1952).
ANBG © M. Fagg, 2006



Source: Australian Plant Image Index (dig.35130).
ANBG © M. Fagg, 2014



Source: Australian Plant Image Index (dig.38581).
ANBG © M. Fagg, 2015



Source: Australian Plant Image Index (dig.38976).
ANBG © M. Fagg, 2015



Source: Australian Plant Image Index (a.100).
ANBG © M. Fagg, 1973



Source: Australian Plant Image Index (a.19345).
ANBG © M. Fagg, 1986



Source: Australian Plant Image Index (a.31305).
ANBG © M. Fagg, 2002




Source: Australian Plant Image Index (dig.15580).
ANBG © M. Fagg, 2009



Source: Australian Plant Image Index (dig.1953).
ANBG © M. Fagg, 2006



Source: Australian Plant Image Index (dig.1954).
ANBG © M. Fagg, 2006



Source: Australian Plant Image Index (dig.35131).
ANBG © M. Fagg, 2014



Source: Australian Plant Image Index (dig.35132).
ANBG © M. Fagg, 2014



Source: Australian Plant Image Index (dig.35133).
ANBG © M. Fagg, 2014



Source: Australian Plant Image Index (dig.38579).
ANBG © M. Fagg, 2015



Source: Australian Plant Image Index (dig.38580).
ANBG © M. Fagg, 2015



Source: Australian Plant Image Index (dig.38582).
ANBG © M. Fagg, 2015



Source: Australian Plant Image Index (dig.38977).
ANBG © M. Fagg, 2015



Source: Australian Plant Image Index (dig.38978).
ANBG © M. Fagg, 2015



Source: Australian Plant Image Index (dig.38979).
ANBG © M. Fagg, 2015



Source: Australian Plant Image Index (dig.19459).
ANBG © M. Fagg, 2011



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.




Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Acacia rigens occurrence map.
Occurrence map generated via Atlas of Living Australia (<https://www.ala.org.au>).

Common Name

Needle Wattle, Nealie, Needle-bush Wattle, Nealia, Niliyah

Family

Fabaceae

Distribution

Mainly found from the Gawler Ra., S.A., E through north-western Vic. to near Dubbo in N.S.W., but with a few scattered occurrences in western S.A., W.A., N.T. and Qld.

Description

Spreading **shrub** 2–3 m high; occasionally **tree** to 6 m high. **Branchlets sericeous** between the yellow ribs. **Phyllodes straight** to shallowly **incurved, terete**, rarely flat, 3–14 cm long, 0.8–1 mm diam., to 2.5 mm wide when flat, **innocuous** or ±coarsely **pungent** with a **straight** or **curved** point, **coriaceous** to subrigid, grey-green, multistriate with often yellowish nerves (some appearing more prominent than others), **appressed-puberulous** or sometimes **glabrous** between the nerves; **gland** basal. Inflorescences **simple**, (1–) 2 (–4) per **axil**; peduncles 2–6 (–8) mm long, **appressed-puberulous**; basal bract persistent; heads **globular**, 20–33-flowered, golden, ±**resinous**. Flowers 5-**merous**; sepals free to 1/2-united. Pods submoniliform, **curved**, 3.5–10 cm long, 2–3 mm wide, firmly **chartaceous** to thinly **coriaceous**, ±**appressed puberulous**. Seeds **longitudinal**, **oblong-elliptic** to narrowly **elliptic**, 3–4.5 mm long, glossy, pale brown; **aril** apical, conical.

Habitat

Commonly in sandy soils but also in red and shaly soils, mostly in mallee scrub.

Specimens

W.A.: 3.2 km E of Boorabbin on Great Eastern Hwy, *B.R.Maslin 1859* (AD, K, MEL, NSW, PERTH). N.T.: 9 km S of Gosses Bluff, *P.K.Latz 17759* (AD, DNA, NT, PERTH).

S.A.: c. 22.8 km SE of Tailem Bend on Dukes Hwy, *N.Hall H80/85* (NSW, PERTH). Qld: Dingwall, 155 km SSE of Charleville, *S.L.Everist 7508* (BRI). N.S.W.: 4.8 km NE of West Wyalong, *R.Coveny 2377* (PERTH). Vic.: 4.8 km SE of Redcliffs, *J.Cullimore 30* (PERTH).

Notes

A widespread and variable species that is often confused with *A. havilandiorum* (S.A., N.S.W., Vic.). Similar in some respects to *A. masliniana* and *A. royei* (W.A.), both of which are readily distinguished by their subulate-pungent phylode apices; also related to *A. argyrotricha* (Qld) and possibly *Acacia* sp. Kalannie North East (B.R.Maslin 7915) (W.A.). One variant of *A. eremophila* was treated as *A. rigens* by B.R.Maslin, *Fl. Central Australia* 130 (1981). A glabrous variant with phylode gland 1.5–2.5 mm above the pulvinus is noted by T.J.Entwistle *et al.*, *Fl. Victoria* 3: 614 (1996) from the Little Desert, Vic.

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

R.S.Cowan

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>

Copyright 2018. All rights reserved.