

WATTLE

Acacias of Australia

Acacia sclerophylla var. *teretiuscula* Maiden & Blakely



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Acacia sclerophylla var. *teretiuscula* occurrence map.
Occurrence map generated via Atlas of Living Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

A poorly collected variety known only from a few localities between Bruce Rock and Lake Grace in south-western W.A.

Description

Branchlets **uncinate-puberulous** (hairs very short, **appressed** to subappressed, abruptly **curved** upwards from base), sometimes almost **glabrous**. New shoots **glabrous** or very sparsely hairy, **resinous**. Stipules **triangular**, c. 0.5 mm long. Phyllodes **linear**, flat to **terete**, 20–45 mm long, 1–1.5 mm wide, **glabrous** or very sparsely **uncinate-puberulous** as on branchlets, with 3 (–5) whitish **longitudinal** nerves. Pods **glabrous** or sparsely **appressed-hairy**.

Habitat

Grows in well-drained, light grey sand or brown clay loam in open shrub mallee woodland.

Specimens

W.A.: between Bruce Rock and Lake Grace [precise locality withheld for conservation reasons], *R.S.Cowan A751 & B.R.Maslin* (BM, BRI, CANB, K, MEL, NSW, PERTH); *loc. id.*, *K.Newbey 1359* (CANB, K, MEL, MO, PERTH).

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

R.S.Cowan, B.R.Maslin

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.