


# WATTLE

## *Acacias of Australia*

### *Acacia* sp. Winged (C.R.Alcock 4936) Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.

### Family

Fabaceae

### Notes

It occurs mainly in coastal areas of S.A. from Venus Bay, western and southern Eyre Penin. and Yorke Penin. *Acacia* sp. Winged (C.R.Alcock 4936) is distinguished most readily from typical *A. anceps* by its generally elliptic, less elongate phyllodes (mostly 2.5–5.5 x 1–3 cm) which have a very reduced pulvinus and which are clearly continuous with the branchlets (that are sometimes narrowly winged) along their adaxial edge, and heads 80–154-flowered. This variant appears to hybridise with the typical *A. anceps* in parts of western Eyre Penin. and also with *A. nematophylla* (see *A. anceps* x *nematophylla*). The type of *A. muelleri*, and probably also those of *A. glaucescens* and *A. pterigoidea*, appear to represent intermediates between the two variants. Further study of this species complex is needed.

This entity is treated as the second variant of *A. anceps* by B.R.Maslin, *loc. cit.*, where it is illustrated by Figs N–O.

### FOA Reference

Flora of Australia Project

### Author

B.R.Maslin, M.O'Leary

This identification key and fact sheets are available as a mobile application:



Australian Government  
Department of the Environment and Energy



Department of  
Biodiversity, Conservation  
and Attractions  
Western Australian Herbarium



Australian  
Biological  
Resources  
Study



POWERED BY  
LUCID®

URL: <https://keys.lucidcentral.org/keys/v3/wattle>

Copyright 2018. All rights reserved.