

# WATTLE

## Acacias of Australia

### *Acacia tetraptera* Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin

Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin

Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
B.R. Maslin

Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.

Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.

Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.

Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.

Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.



Acacia tetraptera occurrence map.  
Occurrence map generated via Atlas of Living  
Australia (<https://www.ala.org.au>).

### Family

Fabaceae

### Distribution

Occurs from near Hyden and Mt Holland ESE to near Grass Patch, south-western W.A.

### Description

Spreading ±open shrub 0.2–0.7 m high; branches slender. Branchlets puberulous to tomentulose. Phyllodes patent, asymmetric, widely ovate to widely depressed-ovate or oblate, (2–) 2.5–4 mm long, 2–4 mm wide, obtuse to acute, apiculate and mucronate, coarsely pungent, thick, smooth and turgid (fresh) but midrib strongly raised when dry, glabrous. Inflorescences simple, 2 per axil; peduncles 4–8 mm long, glabrous; heads globular, 4–5 mm diam., 20–30-flowered, bright light golden to mid-golden. Flowers 5-merous; sepals free. Pods strongly arcuate to subcoiled, subterete, to 2 cm long, c. 2 mm wide, coriaceous to thinly crustaceous, black, glabrous. Seeds longitudinal, oblong, c. 2 mm long, mottled grey and black; aril conical, yellow.

### Habitat

Grows in loam or sand over clayey loam.

### Specimens

W.A.: The Pimple, 40 km E of Hyden, K.J. Atkins 1579 (PERTH); Peak Charles turn-off on Lake King-Kumarl road, B.R. Maslin 5792 (AD, BRI, CANB, G, MEL, MO, NSW, PERTH); 7.5 km E of Grass Patch, D.J. McGillivray 3635 & A.S. George (PERTH).

### Notes

One collection (from 10 km W of Peak Eleanora, M.A. Burgman & S. McNeil 2231, PERTH) appears to represent a possible hybrid with *A. merrallii*.

### FOA Reference


Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

### Author

Minor edits by J.Rogers

B.R. Maslin

This identification key and fact sheets are available as a mobile application:



URL: <https://keys.lucidcentral.org/keys/v3/wattle>  
Copyright 2018. All rights reserved.