

# WATTLE

## Acacias of Australia

### *Acacia vincentii* R.S.Cowan & Maslin



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
R.L. & M.D. Barrett



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
R.L. & M.D. Barrett




Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
R.L. & M.D. Barrett



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
R.L. & M.D. Barrett



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
R.L. & M.D. Barrett



Source: WorldWideWattle ver. 2.  
Published at: [www.worldwidewattle.com](http://www.worldwidewattle.com)  
See illustration.



Acacia vincentii occurrence map.  
Occurrence map generated via Atlas of Living Australia (<https://www.ala.org.au>).

### Family

Fabaceae

### Distribution

Known only by the type collection from Edkins Ra., West Kimberley, W.A.

### Description

Shrub 3 m high with long arching branches. Branchlets terete, commonly glandular-puberulous. Stipules persistent, setaceous, united basally, 2–5 mm long, upcurved. Phyllodes ascending to erect, inequilaterally oblong-oblancoelate, with upper margin conspicuously rounded between apex and gland, (3–)4–6 mm long, 1.5–2 mm wide, l: w = 2–3, with mucronate-pungent apex, coriaceous, ±puberulous, with 2–4 distant slightly raised nerves and occasional anastomoses; gland about middle of adaxial margin. Inflorescences simple, 1 per axil; peduncles 5–10 mm long, puberulous with many hairs gland-tipped; heads obloid, ±5.5 mm diam., 41-flowered; bracteoles long-exserted in buds, spatulate. Flowers 5-merous; sepals 1/2-united. Pods narrowly oblong, with one or both margins crenate, curved, to 5.5 cm long, 5–10 mm wide, crustaceous, obliquely reticulate, densely glandular-puberulous. Seeds (immature) oblique, black, shiny; aril small, apical.

### Habitat

Collected from sandstone plateau.

### Notes

A member of the '*A. deltoidea*' group. Nearest to *A. froggattii* which has globular heads and non-crenate pods with longitudinal seeds; also related to *A. deltoidea* subsp. *deltoidea* which has larger phyllodes of different shape, non-exserted bracteoles, globular heads and non-crenate pods. *Acacia obtriangularis* is also related but easily distinguished by its short-cylindrical spikes.

### FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

### Author

Minor edits by J.Reid

R.S.Cowan

This identification key and fact sheets are available as a mobile application:



Australian Government  
Department of the Environment and Energy



Department of  
Biodiversity, Conservation  
and Attractions  
Western Australian Herbarium



URL: <https://keys.lucidcentral.org/keys/v3/wattle>  
Copyright 2018. All rights reserved.