

WATTLE

Acacias of Australia

Vachellia pachyphloia subsp. *brevipinnula* (Tindale & Kodela) Kodela



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
B.R. Maslin



Source: WorldWideWattle ver. 2.
Published at: www.worldwidewattle.com
See illustration.



Vachellia pachyphloia subsp. *brevipinnula*
occurrence map.
Occurrence map generated via Atlas of Living
Australia (<https://www.ala.org.au>).

Family

Fabaceae

Distribution

Occurs in north W.A., from Prince Regent R. Reserve N to Port Warrender and Cape Londondery, commonly recorded from the Mitchell Plateau.

Description

Branchlets puberulous to pubescent, becoming \pm glabrous on older branchlets. Leaves: main axis puberulous to pubescent, sometimes only sparsely hairy; rachis with jugary glands present at base of all pairs of pinnae or often absent from some pairs of pinnae on leaves with more than 4 pairs of pinnae; pinnae (1–) 2–11 pairs; pinnules up to 54 pairs on pinna-rachis, 3–10 mm long, 1.5–2.5 mm wide. Pods puberulous.

Habitat

Grows in open forest or grassy woodland, in brown or red soils, on basalt, laterite and sandstone, on flat plateau ridges and hillslopes.

Specimens

W.A.: Port Warrender, N Kimberley, *J.S.Beard 6985* (DNA, PERTH); remote weather stn 29 km N of mining camp, Mitchell Plateau, *K.F.Kenneally 8553* (BRI, K, PERTH); Mitchell Plateau area, c. 5 km due SW of Walsh Point, *B.R.Maslin 5131* (NSW, PERTH).

FOA Reference

Data derived from *Flora of Australia* Volumes 11A (2001), 11B (2001) and 12 (1998), products of ABRS, ©Commonwealth of Australia

Author

P.G.Kodela

This identification key and fact sheets are available as a mobile application:



Australian Government
Department of the Environment and Energy



Department of
Biodiversity, Conservation
and Attractions
Western Australian Herbarium



URL: <https://keys.lucidcentral.org/keys/v3/wattle>
Copyright 2018. All rights reserved.